



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

member of the Molgophidæ as one of these connecting types. This localized specialization means that we must look into the Mississippian and the Devonian for the earliest of the Amphibia in North America, as the foot prints which have been discovered in these deposits would indicate.

The discovery of the new temnospondylous form with other facts of the distribution of the Temnospondylia indicates that the order originated in North America. At least the earliest known forms occur in this continent.

The amphibian fauna of Mazon Creek at the present time may be regarded as represented by nine species which are members of four orders and five families. The orders are: Branchiosauria, Microsauria and Temnospondylia. An additional fact of interest is the discovery of osseous branchial arches in an imperfectly preserved specimen; the second species from the Pennsylvanian in which these structures have been seen. This means the presence of a fourth order of Amphibia in the Mazon Creek shales.

ROY L. MOODIE

THE UNIVERSITY OF KANSAS,
January 14, 1910

A FIXING FLUID FOR PLANT TISSUES

My experience with Bouin's fluid as a fixing material for certain plant tissues for cytological work has been so satisfactory that I take this opportunity of recommending it to plant cytologists as one which combines a number of admirable features. It has, of course, been used for a number of years in connection with animal tissues, and especially for studies of spermatogenesis, in which it gives notably clear preparations. I first tried it, along with a number of other solutions, for fixing anthers of *Oenothera*, in 1908. The formula used was as follows:

	Parts
Picric acid, saturated aqueous solution ..	75
Glacial acetic acid	5
Formaline	20

Of course, various modifications of this may be found advantageous for different plant forms.

The time of fixation must be short, otherwise maceration results. It should probably not exceed four to six hours. The time of washing must also be comparatively brief, as long washing causes deterioration and fragmentation of the material. *Oenothera* anthers, after a few hours' immersion in this fluid, frequently acquire a slight pinkish tint, which remains indefinitely after the material has been dehydrated and placed in 70 per cent. alcohol.

This solution seems to be a favorite one for studies on animal spermatogenesis, and I see no reason why it should not become popular also for various purposes in plant cytology. Its obvious advantages are (1) that, unlike osmic solutions, it leaves the tissues clear and transparent, (2) its penetration seems to be very rapid, giving an even and almost perfect fixation of the material, (3) it leaves the cytoplasm and nuclei perfectly colorless, giving particularly clear and brilliant results in staining chromatin and spindles when followed by Heidenhain's iron-hæmatoxylin stain.

R. R. GATES

MISSOURI BOTANICAL GARDEN

THE AMERICAN SOCIETY OF NATURALISTS

The American Society of Naturalists met at the Harvard Medical School, Boston, Mass., on Wednesday, December 29, 1909. There were both morning and afternoon sessions. The program consisted of original papers and demonstrations of studies on evolution, and the meeting proved to be one of the most successful in the history of the society. The variety and importance of the papers read are well shown by the following list of titles:

PAPERS

U. Dahlgren: "Origin of the Electric Tissues in Teleost Fishes" (lantern).

D. T. MacDougal: "Origination of Parasitism in Higher Plants."

F. Boas: "The Influence of Heredity and of the Environment on Man."

E. Brainard: "The Evolution of New Forms in *Viola* through Hybridism."

R. R. Gates: "The Material Basis of Mendelian Phenomena" (lantern).

A. M. Lutz: "The Relation of Chromosome